

Journal of Electron Spectroscopy and Related Phenomena

Journal of Electron Spectroscopy and Related Phenomena 79 (1996) 511-514

A high resolution study of the photoelectron angular distributions and branching ratios for the carbon dioxide molecule in the wavelength region 685-795Å

J. B. West^a, A. C. Parr^b, M. A. Hayes^a, J. L. Dehmer^c, P. M. Dehmer^c, K. Ueda^d, M. R. F. Siggel^c and J. E. Hardis^b

^aDaresbury Laboratory, Warrington WA4 4AD, UK

^bNational Institute of Standards and Technology, Gaithersburg, MD 20899, USA

^cEnvironmental Research Division, Argonne National Laboratory, Argonne, IL 60439, USA

^dResearch Institute for Scientific Measurements, Tohoku University, Sendai 980, Japan



JOURNAL OF ELECTRON SPECTROSCOPY AND RELATED PHENOMENA

EDITORS

C.R. Brundle
(San Jose, CA)

G.E. McGuire
(Research Triangle Park, NC)

J.J. Pireaux
(Namur, Belgium)

EDITORIAL BOARD

P.S. Bagus (San Jose, CA)

H. Bock (Frankfurt)

A.M. Bradshaw (Berlin)

D. Briggs (Middlesbrough, UK)

J. Cazaux (Reims)

M. Delamar (Paris, France)

J.M. Dyke (Southampton, UK)

C.S. Fadley (Berkeley, CA)

S.W. Gaarenstroom (Warren, MI)

A.P. Hitchcock (Hamilton, Ont.)

K. Kimura (Tatsunokuchi)

G.G. Kleiman (Campinas)

A. Kotani (Tokyo)

J.P. Maier (Basel)

N. Mårtensson (Uppsala)

V.I. Nefedov (Moscow)

V.V. Nemoshkaleiko (Kiev)

C. Oshima (Tokyo)

M.N. Piancastelli (Rome, Italy)

C.J. Powell (Gaithersburg, MD)

L. Sabbatini (Bari, Italy)

G.A. Sawatzky (Groningen)

M.P. Seah (Teddington, UK)

K. Siegbahn (Uppsala)

R. Szargan (Leipzig)

S.M. Thurgate (Murdoch, NSW)

S. Tougaard (Odense)

W. von Niessen (Brunswick)

J.H. Weaver (Minneapolis, MN)

SCOPE

This journal publishes experimental and theoretical work in the field of electron spectroscopy and all subjects relevant to it. Subject areas included are physical, organic, inorganic, analytical and theoretical chemistry; surface science and surface analysis; thin film science; semiconductor physics and chemistry; metal surfaces, ceramics, high T_c superconductors, polymers and biomaterials; catalysis; solid state physics; molecular physics. Work can be on free molecules, solids or surfaces. The individual techniques of electron spectroscopy include:

- photoelectron spectroscopy of both outer and inner shells; both UV and X-ray induced
- inverse photoemission
- spin-polarized photoemission
- Auger spectroscopy (including ion neutralization studies)
- synchrotron radiation studies in the area of photoemission and Auger electron emission
- (high resolution) electron energy loss spectroscopy of all types including electronic excitation, high resolution EELS for vibrational studies and $e,2e$ spectroscopy
- electron scattering and resonance electron capture
- electron spectroscopy in conjunction with microscopy
- Penning ionization spectroscopy including tunnelling spectroscopy
- theoretical treatments of the photoemission, Auger, energy loss and Penning ionization processes.

Examples of related topics published in the journal include the theoretical calculations of electronic structure (molecular, solid state band structure, surface electronic structure) and any spectroscopic techniques that are related to the field of electron spectroscopy. These obviously encompass X-ray absorption and fluorescence (including synchrotron studies such as near-edge X-ray absorption fine-structure, NEXAFS), vacuum UV absorption and emission, appearance potential spectroscopy, and mass spectrometric ionization potential measurements. Papers on tunnelling spectroscopy and microscopy are also encouraged.

The journal welcomes papers on instruments and instrumental developments and is one of the major archival sources for work dealing with advances in quantitative analysis by electron spectroscopic techniques, particularly for surfaces and solids. Major topics within this area are ionization cross-sections, angular distributions, mean free path lengths, standards, calibrations, deconvolution procedures, etc.

CATEGORIES OF SCRIPTS

In addition to *Full Papers*, scripts may be classified as Rapid Communications, Review Articles, Data Bank Items and Instrumentation News.

Rapid Communications are short communications of not more than five printed pages including figures and tables. An abstract of not more than 75 words should be included. These communications can be self-contained, complete articles too brief to be classified as full papers, preliminary notes for an ongoing piece of work, extension of previously published work, or comments on published work. The short article should in principle merit accelerated publication. No proofs will be sent to the authors and publication should occur within 3–4 months.

Review Articles will often be by invitation, but contributions are also encouraged. In general the length should not exceed 30 printed pages and authors are recommended to contact one of the Editors or Editorial Board members if in doubt concerning the appropriateness of the subject matter. Authors of review articles may choose between a modest honorarium or 100 extra free reprints: details are available from the Publisher.

Data Bank This category is reserved for electron spectroscopy information in the form of tables or spectra with a minimum of accompanying script. It is really a means of providing data which can be of use to the reader and which would not otherwise be published. Compilations of literature data or references are also occasionally published within this category. There is no page limitation.

Instrumentation News is a category for scripts providing information on instruments or hardware utilized in electron spectroscopy. It is open to commercial instrument manufacturers and individual researchers alike as a place to present description of new instruments, developments of instruments, special components, etc. The scripts will be processed directly by the editors without further refereeing since it is not intended that they be full technical papers; such papers can be submitted within the normal journal categories. This category is intended primarily as a news service to the readers.

A high resolution study of the photoelectron angular distributions and branching ratios for the carbon dioxide molecule in the wavelength region 685–795 Å

J. B. West^a, A. C. Parr^b, M. A. Hayes^a, J. L. Dehmer^c, P. M. Dehmer^c, K. Ueda^d, M. R. F. Siggel^c and J. E. Hardis^b

^aDaresbury Laboratory, Warrington WA4 4AD, UK

^bNational Institute of Standards and Technology, Gaithersburg, MD 20899, USA

^cEnvironmental Research Division, Argonne National Laboratory, Argonne, IL 60439, USA

^dResearch Institute for Scientific Measurements, Tohoku University, Sendai 980, Japan

Measurements of the vibrational branching ratios and photoelectron angular distributions have been made in the regions of the Tanaka-Ogawa, Lindholm and Henning series for the CO₂ molecule. The behaviour of these parameters was found to be sensitive to which particular resonance is excited, with considerable intensity going into vibrational modes other than the symmetric stretch. An initial analysis of some of the data taken is presented.

1. INTRODUCTION

It has been known for some time that the photoionization spectrum of carbon dioxide contains prominent series of autoionizing resonances converging to the A ²Π_u and B ²Σ_u⁺ ionization limits. Although not all the structure has been identified, several vibrational and Rydberg progressions have been and are now known as the Tanaka-Ogawa (TO) series, the Lindholm (L) series and the Henning sharp(s) and diffuse(d) series. Using spectrographic methods, Tanaka and Ogawa[1] identified several Rydberg series converging onto the first eight vibrational levels of the A state, and each of these is split by spin-orbit coupling with the series converging onto the ²Π_{1/2} limit being the stronger. A further progression seen by Tanaka et al[2] was later assigned by Lindholm[3] as a vibrational progression with principal quantum number 3. The other most prominent series were first seen by Henning[4] and converge to the ground vibrational level of the B state of the CO₂⁺ ion.

A full consideration of the symmetries of the series was given recently by Parr et al[5]; in summary, in the case of the TO series some doubt remains but the balance of the experimental and theoretical evidence points to this having an ndδ_g configuration. The Henning sharp and diffuse series are now firmly believed to have ndσ_g and nsσ_g symmetries

respectively.

The study presented here was undertaken to examine the response of angular distribution parameters and branching ratios to complex resonant structure in the photoionization spectrum, with improved electron spectrometer resolution and over a wide wavelength range.

2. EXPERIMENTAL PROCEDURE

This experiment was carried out on a 5 metre normal incidence monochromator fitted to a beamline at the Daresbury SRS, providing a photon flux of ~10¹⁰ photons/sec within a bandpass of 0.1 Å. The light was brought into the experimental chamber by a 2mm diameter capillary light guide, which also served as a pressure differential between this chamber and the ultra high vacuum of the optical monochromator. The electron spectrometer[6] comprised two 100mm radius hemispherical analysers, one rotatable about the incoming light beam as axis and the other fixed. The fixed analyser accepted electrons ejected parallel to the E-vector of the incident radiation, and after calibration procedures were complete the rotatable analyser was set perpendicular to the E-vector. The polarization of the incoming light was measured using a three mirror polarizer with tungsten mesh and plate




Figure 1 The experimental equipment used for the measurements

photodiodes which could be rotated with the rotatable analyser.

Calibration of the energy response of the analysers was done using argon or helium gas, using the known values of the cross section[7] and β -parameters[8]. Throughout the whole wavelength region studied here, measurements were taken at 0.1\AA intervals, and the β -parameters and branching ratios calculated from the measurements made at the two angles using the well known expression

$$\frac{d\sigma}{d\Omega} = \frac{\sigma}{4\pi} \left[1 + \frac{\beta}{4} (3p \cos 2\theta + 1) \right]$$

where θ is the angle between the major polarization axis and the ejected electron, p is the polarization of the incoming light, and σ is the partial cross section for the channel corresponding to the photoelectron being detected. In the case of this experiment, therefore, measurements were made simultaneously at $\theta = 0^\circ$ and $\theta = 90^\circ$. Altogether ~ 1500 data sets were taken in the wavelength region 650\AA to 890\AA . The data analysis

was carried out by first correcting the raw data for the differing detection efficiencies of the two electron energy analysers. After normalization for detection efficiencies the data were fitted to a series of gaussian profiles using a least squares procedure, using as a basis the known levels of CO_2^+ identified by Chambaud et al[9]. The data were of sufficient quality to give reliable intensities for the vibrational components down to the 5% level.

3. RESULTS

With the large amount of data and the level of detail available to us, in this initial analysis we have concentrated on the response of the symmetric stretch vibrational members of the CO_2^+ ion to resonant structures. For the purposes of convenience and brevity, we have followed the notation adopted by McCulloh[10] to label the series members; for example $A_{5_2}(\text{TO})$ means the member of the Tanaka-Ogawa series for which the running number is 5 and the

vibrational quantum number is 2. Bearing in mind that this series is now believed to have $nd\delta_g$ symmetry, the principal quantum number should be 4, ie one less than shown on the graphs. In the case of the Henning series, though using McCulloh's notation, we follow the reassignment given by Fridh et al[11], and the running numbers shown are the principal quantum numbers. This results in our labelling of the resonances being different from that in McCulloh's table.




Figure 2. Branching ratios for the (000) vibrational members of CO_2^+ for the A4(TO) series.

We show in figure 2 the vibrational branching ratios for the first eight members of the symmetric stretch vibrational members in the CO_2^+ ion ground state, for the (TO) A_4 series of resonances. Although we can resolve the spin orbit split members of these series,




Figure 3. Branching ratios for the (000) vibrational members of CO_2^+ for the Henning sharp series. Note that in this figure the resonances have been numbered to avoid overcrowding on the diagram for the higher Rydberg members. The actual wavelengths are readily available from the literature; see, for example, reference 10.

the $^2\Pi_{1/2}$ components, at the shorter wavelength of the pair, have the larger intensities and are the ones to which we refer on the figures.

The differing response of the CO_2^+ ion vibrational members to the resonances is obvious; although the (000) member always has the largest intensity the sum of the branching ratios at some resonances falls well below 0.5, indicating that a substantial intensity goes into the bending and anti-symmetric stretch vibrations. This is less true for the Henning series, for which the corresponding branching ratio data are shown in figure 3; in this case it can be seen that the (000) member tends to be larger, in comparison with the A4(TO) series.

With our experimental resolution of 40 meV, much of the weaker structure in the photoelectron spectrum is not resolved, nor is the spin-orbit splitting of the $A\ ^2\Pi_u$ state. This limits us in the number of vibrational levels for which we can provide reliable angular distribution and branching ratio values, and is reflected in the error bars shown in the data, derived mainly from the quality of the least squares fit to the electron spectrum. However, in a separate experiment[5] using jet-cooled CO_2 and a spectrometer resolution of 17 meV, more structure was resolved. The results in figure 4 were taken at the peak of the $A5_2$, ie $m = 5$, (200) member of the TO series. This experiment was undertaken to show the selectivity in the autoionization of this state, and as can be seen it decays primarily to the $^2P_{1/2}$ level, consistent with the finding that the (TO) series converging to the $^2P_{1/2}$ limits are much more intense than those converging to the $^2P_{3/2}$ limits. From symmetry considerations[5] these data indicate that the (TO) series should be described by $\Omega_c\omega$ coupling. The known energy levels of members of the (v_100) , (v_101) and (v_102) progressions are shown as solid lines; the positions of the higher members were estimated by linear or quadratic extrapolation of the known levels. Also the Renner-Teller split (020) band is clearly seen.

The analysis and interpretation of these results is currently under way and will be the subject of a forthcoming publication.

4. ACKNOWLEDGMENTS

This work was supported by the U.S. Department of Energy, Office of Energy Research, under Contract




Figure 4. Photoelectron spectrum taken at the $A5_2$, $m=5$, (200) member of the (TO) series, at an analyser angle of 0° . The off-resonance contribution has been subtracted.

No. W-31-109-ENG-38 and by the Engineering and Physical Sciences Research Council, UK.

5. REFERENCES

1. Y. Tanaka and M. Ogawa, *Can. J. Phys.* 40 (1962) 879
2. Y. Tanaka, A.S. Jursa and F.J. LeBlanc, *J. Chem. Phys.* 32 (1960) 1199
3. E. Lindholm, *Arkiv Fysik*, 40 (1969) 129
4. H. J. Henning, *Ann. Phys. (Leipzig)* 13 (1932) 599
5. A.C. Parr, P.M. Dehmer, J.L. Dehmer, K. Ueda, J.B. West, M.R.F. Siggel and M.A. Hayes, *J. Chem. Phys.* 100 (1994) 8768
6. A.C. Parr, S.H. Southworth, J.L. Dehmer and D.M.P. Holland, *Nucl. Instrum. Methods*, 222 (1984) 221
7. G.V. Marr and J. B. West, *At. Data and Nucl. Data Tables*, 18 (1976) 497
8. D. M. P. Holland, A. C. Parr, D. L. Ederer, J. L. Dehmer and J. B. West, *Nucl. Instrum. Methods*, 195 (1982) 331
9. G. Chambaud, W. Gabriel, P. Rosmus and J. Rostas, *J. Phys. Chem.* 96 (1992) 3285
10. K. E. McCulloh, *J. Chem. Phys.* 59 (1973) 4250
11. C. Fridh, L. Åsbrink and E. Lindholm, *Chem. Phys.* 27 (1978) 169

INSTRUCTIONS TO CONTRIBUTORS

Copies of the Guide for Authors are available from the Editors and the Publisher, and have been printed in the journal as *J. Electron Spectrosc. Relat. Phenom.*, 67 (1994) 521–527.

SUBMISSION OF PAPERS AND COMPUSCRIPTS (IN ENGLISH)

Hardcopy scripts should be submitted in triplicate to one of the Editors. **Instructions for the preparation of electronic scripts and illustrations are available from the Publisher and the Editors.** In order to speed up the publishing process, papers from Europe and the former Soviet Union should preferably be sent to Professor J.J. Pireaux for further handling. Papers from Asia and Australasia should be submitted to Dr C.R. Brundle. All other papers should be sent to either Dr C.R. Brundle or Dr G.E. McGuire.

C.R. Brundle, IBM Almaden Research Center, K31/802, 650 Harry Road, San Jose, CA 95120-6099, USA.

G.E. McGuire, Microelectronics Center of North Carolina, P.O. Box 12889, Research Triangle Park, NC 27709-2889, USA.

J.J. Pireaux, Lab. Interdisciplinaire de Spectroscopie Electronique, Facultés Universitaires Notre Dame de la Paix, rue de Bruxelles 61, B-5000 Namur, Belgium.

Submission of an article is understood to imply that the article is original and unpublished and is not being considered for publication elsewhere.

SCRIPTS

Scripts should be typed (double-spaced) on one side only of consecutively numbered pages. Illustrations should be clearly numbered. Line drawings should be drawn in Indian ink on drawing or tracing paper and marked with standard symbols. Authors are reminded that papers appear in two columns, each 7.5 cm by 20.5 cm (total type area 16 cm by 20.5 cm), and illustrations should be produced to suit the format of the journal. Photographs should be submitted as clear black and white glossy prints. The Publisher will return to authors the original figures of a printed article if requested at the time of submission. Figure legends and tables should be typed together on separate pages. The reference list should appear at the end of the paper and should be numbered in order of the appearance of the references in the text (not arranged alphabetically). Reference numbers in text should appear on the line between square brackets. Abbreviations of journal titles should conform to those adopted by the Chemical Abstracts Service (Bibliographic Guide for Editors and Authors, The American Chemical Society, Washington, DC, 1974). The recommended form for references to journal papers and books is as follows:

[1] E.O.F. Zdansky, M. Ohno and N. Mårtensson, *J. Electron Spectrosc. Relat. Phenom.*, 67 (1994) 141.

[2] G.N. Belozerski, Mössbauer studies of surface layers, *Studies in Physical and Theoretical Chemistry*, Vol. 81, Elsevier, Amsterdam, 1993.

[3] R.D. Thomas, in E. Buncel and J.R. Jones (Eds.), *Isotopes in the Physical and Biomedical Sciences*, Vol. 2, Elsevier, Amsterdam, 1991, Chapter 7.

[4] G. Gensterblum, J.-J. Pireaux, P.A. Thiry, R. Caudano, Ph. Lambin and A.A. Lucas, in M. Rocca, G. Bracco and U. Valbusa (Eds.), *Vibrations at Surfaces 1993*, Proc. 7th International Conference on Vibrations at Surfaces, Santa Margherita Ligure, Genoa, Italy, 14–17 June 1993, Elsevier, Amsterdam, 1993, p. 835.

For multi-author references, all authors must be named, and initials given, in the reference list although the use of, for example, Smith et al. is desirable in the text. Authors are requested not to use any underlining in the reference list.

REPRINTS

There are no page charges and fifty reprints will be supplied free of charge. Additional reprints (minimum 100) can be ordered at quoted prices. They must be ordered on order forms which are sent together with the proofs.

PUBLICATION AND SUBSCRIPTIONS

The *Journal of Electron Spectroscopy and Related Phenomena* (ISSN 0368-2048). For 1996, volumes 75–81 are scheduled for publication. Subscription prices are available upon request from the Publisher. Subscriptions are accepted on a prepaid basis only and are entered on a calendar year basis. Issues are sent by surface mail except to the following countries where air delivery via SAL (Surface Air Lift) is ensured: Argentina, Australia, Brazil, Canada, Hong Kong, India, Israel, Japan, Malaysia, Mexico, New Zealand, Pakistan, PR China, Singapore, South Africa, South Korea, Taiwan, Thailand, USA. For all other countries airmail rates are available upon request. Claims for missing issues must be made within six months of our publication (mailing) date.

Please address all your requests regarding orders and subscription queries to Elsevier Science B.V., Journal Department, P.O. Box 211, 1000 AE Amsterdam, The Netherlands. Tel: +31 20 580 3642; Fax: +31 20 580 3598.

ADVERTISEMENTS: for rates apply to the Publisher.

© 1996, ELSEVIER SCIENCE B.V. ALL RIGHTS RESERVED.

0368-2048/96/\$15.00

No part of this publication may be reproduced, stored in a retrieval system or transmitted in any form or by any means, electronic, mechanical, photocopying, recording or otherwise, without the prior written permission of the Publisher, Elsevier Science B.V., Copyright and Permissions Department, P.O. Box 521, 1000 AM Amsterdam, The Netherlands.

Upon acceptance of an article by the journal, the author(s) will be asked to transfer copyright of the article to the Publisher. The transfer will ensure the widest possible dissemination of information.

Special regulations for readers in the USA – This journal has been registered with the Copyright Clearance Center, Inc. Consent is given for copying of articles for personal or internal use, or for the personal use of specific clients. This consent is given on the condition that the copier pay through the Center the per-copy fee for copying beyond that permitted by Sections 107 or 108 of the US Copyright Law. The per-copy fee is stated in the code-line at the bottom of the first page of each article. The appropriate fee, together with a copy of the first page of the article, should be forwarded to the Copyright Clearance Center, Inc., 222 Rosewood Drive, Danvers, MA 01923, USA. If no code-line appears, broad consent to copy has not been given and permission to copy must be obtained directly from the author(s). The fees indicated on the first page of an article in this issue will apply retroactively to all articles published in the journal, regardless of the year of publication. This consent does not extend to other kinds of copying, such as for general distribution, resale, advertising and promotion purposes, or for creating new collective works. Special written permission must be obtained from the Publisher for such copying.

No responsibility is assumed by the Publisher for any injury and/or damage to persons or property as a matter of products liability, negligence or otherwise, or from any use or operation of any methods, products, instructions or ideas contained in the material herein. Although all advertising material is expected to conform to ethical (medical) standards, inclusion in this publication does not constitute a guarantee or endorsement of the quality or value of such product or of the claims made of it by its manufacturer.

This issue is printed on acid-free paper.

PRINTED IN THE NETHERLANDS